Smell-e Technology: Bridging the gap between virtual and real-life food responses using an

immersive multisensory VR food environment

Nynke van der Laan ^a L.N.vdLaan@tilburguniversity.edu

Sanne Boesveldt ^b ^b Wageningen University & Research Sensory Science & Eating Behaviour chairgroup - Division of Human Nutrition & Health <u>sanne.boesveldt@wur.nl</u>

> Matti Vuorre[°] [°]Tilburg University Department of Social Psychology <u>M.J.Vuorre@tilburguniversity.edu</u>

Tessa M. van Leeuwen ^{a,d} <u>T.M.vanLeeuwen@tilburguniversity.edu</u> ^d Radboud University Donders Institute for Brain, Cognition, and Behaviour

> Kim Verboon ^a <u>k.g.verboon@tilburguniversity.edu</u>

Travis Masterson ^e ^d The Pennsylvania State University Department of Nutritional Sciences <u>travis.d.masterson@psu.edu</u>

Alexander Klippel^f ^e Wageningen University & Research Cultural Geography Research Group & WANDER XR Experience Lab <u>Alexander.klippel@wur.nl</u>

> Rachelle de Vries ^{a,1} (Corresponding Author) ^a Tilburg University Department of Communication & Cognition P.O. Box 90153 5000 LE Tilburg (The Netherlands) <u>rachhdevries@gmail.com</u>

¹ Present address: Unilever Foods Innovation Centre, Science & Technology Future Flavour Team, Bronland 14, 6708 WH Wageningen (The Netherlands)

Abstract

Immersive Virtual Reality (VR) technologies such as virtual supermarkets are an emerging medium to model individuals' eating behaviour. However, existing VR environments elicit weaker responses to food (i.e., craving and salivation) than in real-life, limiting their validity as research tools. We developed an immersive multisensory VR food environment – with both visual and olfactory (smell) cues – and investigated whether it could bridge this gap in food responses, and whether effects may be mediated by an enhanced sense of presence. In a within-subjects lab-based experiment, participants (N = 70) were exposed to food and non-food cues in either a unisensory "vision only" VR condition, a multisensory "vision + olfaction" VR condition, or a real-life setting with a matched physical set-up. Foodspecific craving and salivation were measured in all six conditions. Results showed that foodinduced craving was weaker in all virtual conditions versus real-life. Salivary responses to food were also lower in unisensory VR exposure versus real-life. Compared to unisensory VR exposure, multisensory VR exposure led to a directional improvement in craving, higher salivary food responses after adjusting for hunger, and enhanced perceptions of presence and mental imagery. While we could not conclude equivalence between multisensory VR and real-life settings, the latter did not differ on salivary responses either. In conclusion, an immersive multisensory VR food environment with olfactory cues can credibly model craving responses, albeit to a weaker degree than in real-life. The added value of this technology may lie in enhancing conceptual mediators and approximating real-life salivation to food.

Keywords: Immersive VR; Virtual supermarket; Olfaction; Food cue exposure; Craving; Salivation; Presence

1. Introduction

Many individuals struggle with making healthier food choices, as evidenced by the growing prevalence of obesity worldwide (World Health Organization, 2020). This is in part because the majority of our food decisions are made in sensory-rich "obesogenic" food environments (e.g., supermarkets) that promote the purchase and consumption of unhealthy foods (Poelman *et al.*, 2021; Swinburn *et al.*, 2011). Therefore, in order to encourage healthier food decisions and ultimately improve diets, we urgently need to better understand the mechanisms driving food choice within such contexts. In this paper, we focus on improving the methods for studying eating behaviour in a lab environment.

Immersive Virtual Reality (VR) technology has recently surfaced as a promising medium to study food decision making in naturalistic food environments (Blom *et al.*, 2021; Xu *et al.*, 2021). Immersive virtual environments are computer-generated three dimensional models that participants can experience and interact with intuitively in real time. Through naturalistic interaction, enabled by a head-mounted display and hand-held controllers that provide a high level of sensory immersion (Slater, 2018), individuals can experience a strong sense of (spatial) presence — i.e., the feeling of being in the virtual environment rather than the physical one (e.g., Lombard and Ditton, 1997; Slater & Wilbur, 1997). Such virtual environments, for example, virtual supermarkets, buffets, and other food environments, have opened up new opportunities for eating behaviour research, allowing researchers to collect food decision-making data in a tightly controlled yet realistic environment, at relatively low cost and with a high degree of flexibility (Xu *et al.*, 2021).

Though previous studies with low-immersive VR (e.g., desktop VR; Waterlander *et al.*, 2015) and semi-immersive setups (e.g., employing multiple screens; van Herpen *et al.*, 2016) have shown that consumer behaviour in virtual environments resembles that in real-life moderately to well, systematic comparisons involving high-immersive VR environments are

lacking (for some exceptions see Cheah *et al.*, 2020 and Long *et al.*, 2023). It is therefore relevant to address this gap (further) because of the expectation that behaviour in highly immersive environments would align more closely with actual behaviour in the real-world due to the stronger level of immersion provided by the medium.

An important prerequisite for immersive technologies to elicit behaviour similar to those in real-world circumstances, and thus serve as accurate tools for measuring eating behaviour, is that basic food cue responses (FCRs) to virtual foods should be similar to those elicited by the same foods in physical (i.e., real-life) settings. FCRs, which include psychological responses (e.g., craving) and physiological responses (e.g., salivation), are necessary to prepare the body for ingestion and represent behaviorally-relevant markers of the cephalic phase of the digestive process. The manifestation of such responses has been found to be a predictor of food choice and food intake (Hill, 2007; Kanoski & Boutelle, 2022; Nederkoorn *et al.*, 2000). However, recent evidence from a highly immersive VR study, in which participants could interact with a virtual versus real-life version of a food, suggests that important psychological FCRs (i.e., craving) tend to be weaker for virtual compared to real-life foods (van der Waal *et al.*, 2021). Similarly, fundamental physiological FCRs (i.e., salivation) to virtual food cues show a much larger departure from reality (van der Waal *et al.*, 2021). As such, weaker psychological and physiological FCRs currently limit the utility of highly immersive virtual environments for food decision-making research.

We argue that the discrepancy in FCRs between (immersive) virtual and real-life foods is because existing VR technologies only serve the visual sense. Though the visual sense plays an important role in food choice, food choice is a multisensory phenomenon that is also driven by smell, taste, texture, and sound input (Motoki & Togawa, 2022). Here, we take the challenging step of developing and testing a novel immersive multisensory VR food environment – with both visual and olfactory (smell) cues – to help bridge this gap. While the

inclusion of visual and auditory senses into VR is widespread, the potential of the sense of smell remains largely untapped (Neo *et al.*, 2021; Xu *et al.*, 2021). The few studies that have included smell in a digital environment have focused solely on the development of the technology (e.g., usability and feasibility tests; Liu *et al.*, 2023; Niedenthal *et al.*, 2023), or they have applied it in a non-food domain such as for the 'gamification' of olfactory cognition testing and training (Andonova *et al.*, 2023; Dozio *et al.*, 2021; Olofsson *et al.*, 2017). However, smell is an established determinant of flavor perception and food choice (Boesveldt & de Graaf, 2017). Indeed, food odours are known to trigger specific appetite (Ramaekers *et al.*, 2014), can enhance the effect of visual food stimuli on self-reported craving (Wolz *et al.*, 2017) and salivation levels (Krishna *et al.*, 2014), and potentially influence food decision-making processes (Morquecho-Campos *et al.*, 2022; Yang *et al.*, 2023).

A mechanism by which an immersive multisensory VR food environment with both visual and olfactory cues may elicit responses more similar to those in real-life is the increased levels of spatial presence within the virtual environment. We propose that the additional olfactory cues enhance the sense of presence, which in turn triggers more realistic responses to the environment. An environment that closely resembles real-life sensory input is expected to give users a greater feeling of actually "being there". Indeed, it has been shown that more sensory-rich (i.e., where more senses are stimulated) VR environments elicit stronger levels of (tele) presence (Goncalves *et al.*, 2020; Galace, 2012), though research on the specific addition of only smell is limited. The few studies that looked at the isolated effect of (environment-congruent) smell exposure added to a VR application generally found positive effects on levels of spatial presence (Archer *et al.*, 2022; Baus *et al.*, 2017; Baus *et al.*, 2022; Brengman *et al.*, 2022; Persky & Dolwick, 2020; Munyan III *et al.*, 2016), though some null findings have also emerged (e.g., Baus *et al.*, 2018).

Several studies have shown that spatial presence is a central conceptual mediator of many VR applications (Barranco Merino *et al.*, 2023; Slater & Wilbur, 1997). Intuitively, when people have a stronger sense of actually being in the virtual environment, they are more likely to respond to stimuli in this environment as they would in physical (i.e., real-life) settings. Indeed, it has been shown that high presence in a virtual environment leads to behaviours more similar to those in real-life circumstances. For instance, in a non-immersive virtual supermarket, participants who reported higher levels of presence had greater similarity in product purchases between virtual and real-world (i.e., non-lab) environments (Waterlander *et al.*, 2015). Even though some evidence exists for the effect of olfaction on spatial presence and the role of spatial presence in eliciting real-life purchasing behaviours, the mediating effect of spatial presence is rarely investigated in food decision-making contexts. Understanding this is crucial, as presence-enhancing factors like smell can improve the ecological validity of virtual supermarkets, and help provide a powerful tool for studying food decision-making. However, the merits of an immersive multisensory VR food environment in this context have yet to be fully assessed.

Therefore, the key objective of the present research was to systematically examine the validity of an immersive multisensory VR food environment, which includes visual and olfactory food cues, for modelling FCRs as measured in a real-life lab setting. Here we address the following research questions: 1) To what extent do individuals' psychological (i.e., craving) and physiological (i.e., salivary) FCRs differ between a unisensory (vision only) VR environment versus a multisensory (vision + olfaction) VR environment versus a real-life setting? 2) What psychological mechanisms underlie FCR-enhancing effects of the multisensory VR environment?

We hypothesized that food cues would lead to stronger craving and salivation responses compared to non-food cues in general (H_{1A}), but that these food-specific cue responses (FCRs)

are smaller in the Unisensory VR environment compared to both Multisensory VR (H_{1B}) and Real-life exposure (H_{1C}). We did not expect a difference in FCR profiles in the latter two exposure modes, in that FCRs would be similar across Multisensory VR and Real-life environments (H_{1D}). Finally, we expected that the difference in FCRs (Δ FCRs) between Multisensory VR and Unisensory VR exposure is (partially) mediated through an enhanced sense of presence (H_2). That is, Multisensory VR exposure will lead to higher levels of presence than Unisensory VR exposure, and presence perceptions will positively correlate with FCRs.

2. Methodology

2.1 Design

This lab-based experiment had a 3 (*Exposure Mode*: Real-life versus Unisensory VR versus Multisensory VR) by 2 (*Stimulus Type*: Non-food versus Food) within-subjects design. We chose a within-subjects design in light of high individual variability in salivary flow rates (Dawes, 1987; Ship *et al.*, 1991). Participants visited the lab twice for screening and the test session, respectively, with a washout period of at least one day. Test sessions were planned between 9:00 and 17:00, since this interval spans typical mealtimes and circadian rhythms governing salivary flow tend to peak around this period (Dawes, 1975; Dawes, 1996).

During test sessions, participants performed all six experimental conditions of a cue exposure task (cf. section 2.4.1) in a hungry state. Hunger was expected to trigger the strongest (psychological and physiological) responding to food cues (Brunstrom *et al.*, 2004; Burgess *et al.*, 2016; Loeber *et al.*, 2013; Steel *et al.*, 2006), as well as better differentiate food cue responses between virtual and real-life conditions (cf. van der Waal *et al.*, 2021). Notably, we pseudorandomized the order of exposure modes and counterbalanced the presentation of stimulus types across participants (Figure 1). The study design, hypotheses,

and analytical plan were pre-registered and are available with study data on the Open Science Framework (Project URL: <u>osf.io/n8gm3/</u>).



Figure 1. Randomization of participants into the different orders of exposure modes. The order in which stimulus types (within an exposure mode) was presented was fixed per participant and counterbalanced across our sample.

2.2 Participants

A total of 70 participants (59 % F; Age = 20.71 (SD = 2.55) years; BMI = 22.39 (2.67) kg/m2) were included in the study. All participants were English-speaking university students from various educational backgrounds (i.e. under- and postgraduate). Participant recruitment was achieved through advertisement of study posters and flyers on university buildings, social media platforms and the participant pool of the university department. Individuals were included in the study if they were healthy at the time of study (self-reported) and had a normal olfactory ability, as assessed using age-specific cut-offs from the 16-item Sniffin' Sticks identification test (Hummel *et al.*, 2007). Individuals were not allowed to participate if they reported a restriction or aversion to test stimuli (cf. section 2.4.3), a presence or history of eating disorder(s), neurological, and/or olfactory disorder(s), identified as being a habitual smoker, or were pregnant and/or lactating at the time of the study. All participants provided written informed consent prior to testing and were compensated with

study credits. This research was approved by the Ethical Review Board of the Tilburg School of Humanities and Digital Sciences (Tilburg University; file number: REDC2023.62).

2.3 Procedure

Experimenters communicated the cover story that the study aimed to investigate effects of VR exposure on cognitive performance. Interested individuals were first screened on their eligibility in a first lab (screening) session, in which their olfactory ability was examined. Eligible participants then received instructions to adhere to prior to the second lab session (test session), including a restriction on using fragranced products (e.g., perfume, chewing gum) and a directive to drink enough water (as per usual consumption) on the day itself. Importantly, to experimentally manipulate hunger states, we also instructed participants to not consume anything (except water) for at least three hours prior to testing (cf. van der Waal *et al.*, 2021). Participants were emailed a reminder for these instructions 24 hours before their test session.

At the onset of a test session, participants filled in a questionnaire on their current hunger level, pre-test adherence checks (e.g., time of last meal), and demographic (e.g., *Gender, BMI*) and control measures (e.g., *familiarity with VR, trait sensory imagery*). Next, they were asked to rinse their mouths with distilled water, which was done to help participants practice the spitting method (cf. section 2.5.1), as well as stabilize salivary levels prior to the cue exposure task (section 2.4.1). To familiarize participants with the VR equipment and the virtual supermarket, they first engaged in a short practice round where they were instructed to pick-up an object in a separate section of the virtual supermarket. Next, they had to complete a cue exposure task in each of six experimental conditions, in a randomized order. Participants had a two-minute break in-between experimental conditions, in which they (re)rinsed their mouths and performed a timed (non-food) distractor task that was related to the cover story (e.g., memory and spatial rotation tests). Attentional checks and

honesty reminders were administered at main procedural checkpoints (e.g., before craving ratings following cue exposure in each condition). Upon finishing the cue exposure task in the last (sixth) condition, participants were debriefed and compensated.

2.4 Apparatus and Stimuli

2.4.1 Cue exposure paradigm. Individuals were instructed to "*imagine you are* grocery shopping in a supermarket, when you come across a promotional stand for a new product". They were then told to explore a sample of the new product up-close, by picking it up from its display (i.e., bowl) and to interact with the sample as they wished to for one minute. They were also informed not to explore any other objects or navigate away from the promotional stand during this time.

For each condition, participants had to interact with the (non-food or food) product for a duration of one minute. Prior to the interaction with the product, individuals were required to swallow immediately before they picked-up the sample. Every 30 seconds after (i.e., twice in total), they would then receive a verbal cue from the experimenter to drop the sample back into its bowl and to spit their saliva into an empty pre-weighted cup. Importantly, in line with the spitting method of Navazesh (1993), individuals were explicitly told not to swallow their saliva – but to let it pool or collect in their mouths – during each 30second interval. After the timed cue exposure, they answered questionnaires (e.g., craving, state gustatory mental imagery, and presence in virtual conditions only) related to their interaction with the product on a provided tablet. This procedure was repeated for each experimental condition. We measured salivary volume as our physiological FCR, while reported craving was our psychological FCR outcome.

2.4.2 Exposure mode. The immediate (visual) environment was standardized as much as possible across exposure modes. In all conditions, participants stood in front of a promotional stand that consisted of a table with a packaged product and unpackaged sample

(within a bowl) on display, as well as an accompanying signboard that stated "New Product – Try Me!". We opted for participants to interact with the unpackaged sample within the bowl to stimulate consumption beliefs (or imagery) in food cue conditions, which may be a prerequisite for salivary responses to occur (Spence, 2011).

In the *Real-life* exposure condition, the promotional stand (i.e., table, promotional sign, packaged product and unpackaged sample within a bowl) was recreated in a separate lab room, with actual (non-food and food) stimuli that participants could interact with (Figure 2).

In the Unisensory VR (i.e., vision only) condition, this promotional stand was placed in a virtual supermarket (VirtuMart; Blom *et al.*, 2021; van der Laan *et al.*, 2022; **Figure 2)**. The VirtuMart is a virtual supermarket modelled in Blender and implemented in Unity. The layout and products of this supermarket were modelled to mimic one of the most well-known Dutch supermarket chains. The assortment of VirtuMart comprises 240 products across twenty product categories, including bread, desserts, meat, fruit, and vegetables. An immersive experience of the virtual supermarket was delivered by using an HTC Vive headmounted display; two hand-held controllers allowed for picking up products using virtual hands. The promotional stand was placed in front of the virtual bread aisle.

The *Multisensory VR* (i.e., vision + olfaction) exposure mode was identical to the unisensory VR condition described above, except that virtual products were further accompanied by olfactory cues. The multisensory VR infrastructure entailed an additional portable olfactometer (*Sniff-O* device developed by *CyNexo srl*) that was connected to an air pressure generator (operating at 3.5 to 4 bar). *Sniff-O* had six odour channels that were equally divided among (non-food and food) stimulus types. Each odour channel had a respective odour jar containing a scented cotton ball and downstream tubing, which ran from the jar and converged into a user manifold that had two main output tubes. The user manifold was fastened to a body harness that was placed at a comfortable distance (i.e., on the center of

the chest bone) and directed odours towards the nose of the participant – adjusting the placement of the manifold per individual as needed (Figure 2). This "free-hanging" construction allowed for hand controllers to be used during the cue exposure task.

Sniff-O interfaced with the VirtuMart Unity platform using an Arduinobased architecture. Two odour channels were operational (i.e., one per stimulus type). Virtual "triggers" were programmed into the VR Unity platform, such that the appropriate odour channel of Sniff-O automatically opened when the corresponding virtual product was pickedup and immediately closed when the object was dropped. In addition, when an odour channel closed, an odourless clean-air channel was simultaneously activated, which produced a continuous flow of air and prevented the cross-contamination of odours within main output tubes. During odour administration, a total flow rate of 3.5 standard liters/minute was used (similar to Albayay et al., 2022): The clean-air (constant flow) channel was delivered at a flow rate of 2 standard liters/minute and odour channels were delivered at a flow rate of 1.5 standard liters/minute. These values were informed by pilot tests and decided on the basis of a stable perceived odour intensity throughout the (one-minute) cue exposure time (see odor pilot in next section). Flow rates were calibrated at the beginning of each test day. Moreover, scented cotton balls were replaced regularly (i.e., every 2-3 test days) to maintain a consistent odour intensity. Finally, in line with best practices (e.g., Lundstrom et al., 2010), odour tubes were "flushed" at the end of each test week, by running clean pressurized air throughout the entire tubing system.

2.4.3 Stimulus type. *Non-food* stimuli encompassed small-to-medium sized scented wood chips (Whiskey wood chips; *Weber*). This product was chosen because wood was previously validated as a suitable non-food reference (cf. van der Waal *et al.*, 2021), contextually appropriate for the supermarket scenario, and matched prominent visual characteristics (e.g., shape, size, and colour) of the food stimuli.



Figure 2. Illustration of the Real-life, Unisensory VR, and Multisensory VR exposure modes, across stimulus types (Non-food versus Food). In the Multisensory VR condition (last panel), visual cues were additionally accompanied by olfactory cues that were administered by a novel portable olfactometer device (*Sniff-O*). Odours were "triggered" as soon as a participant picked up the corresponding object (i.e., wood versus chocolate) in the virtual food environment.

Food stimuli consisted of unwrapped (pure) milk chocolate pieces. Chocolate is

widely recognized as a rewarding food and has been demonstrated to be a potent stimulus for

inducing strong appetitive responses (Kemps & Tiggemann, 2009; Proserpio et al., 2017).

For virtual conditions, the same three-dimensional (unwrapped) chocolate model from

van der Waal *et al.*, (2021) was used. All remaining virtual products (e.g., packaged chocolate, and (un)packaged wood) were developed from scratch using Blender software and modelled as closely as possible to real-life variants. For both wood (AllSense-Voit Aroma Factory No. 821; 1.2% in propylene glycol) and chocolate (IFF SC048015; 5% in propylene glycol) odour counterparts, odour solutions were first created and a small volume of each was then pipetted into cotton balls for insertion into *Sniff-O* (3 and 2 milliliters for wood and chocolate, respectively). We piloted a set of odour dilutions in a separate student sample (N=35) beforehand to arrive at the selected concentrations (see Table S1), ensuring a similar rated liking and medium-high perceived intensities (i.e., 55-75 mm on a 100 mm VAS) among non-food and food stimuli. Furthermore, final odours were consistently (i.e., more than 80% of the time) correctly categorized as (non)food and matched with their associated object significantly more frequently than chance level (Table S1).

2.5 Measurements

2.5.1 Primary outcome variables. To assess psychological cue responses, we asked individuals to rate their subjective *craving* for chocolate in all six experimental conditions. *Craving* was determined by the statement "*How much do you desire to eat chocolate at this moment*" rated on a 100 mm VAS anchored from "Not At All" to "Very Much" (Hill, 2007; van der Waal *et al.*, 2021). We then took the difference in *craving* between food and non-food stimuli to accurately quantify psychological FCRs in each exposure mode.

We measured individuals' physiological cue responses by collecting whole-mouth *salivary volume* using the spitting method (cf. Navazesh, 1993) and weighing the difference (in grams; *g*) of a salivary cup before and after cue exposure in each condition (Morquecho-Campos *et al.*, 2019; van der Waal *et al.*, 2021). Similarly, we used the difference in *salivary*

volume between food and non-food stimuli in each exposure mode as a proxy for physiological FCRs.

2.5.2 Process indicators. *Presence* in virtual conditions was examined using the 14-item Igroup Presence Questionnaire (Schubert *et al.*, 2001), which focuses on the concept of "being there" or one's degree of engagement with the virtual as opposed to real (physical) world (Grassini & Laumann, 2020). The questionnaire is divided into three subscales: spatial presence (e.g., "Somehow I felt that the virtual world surrounded me"), involvement (e.g., "I was not aware of my real environment"), and experienced realism ("The virtual world seemed more realistic than the real world"), all rated on a five-point Likert scale. The scale had a good internal reliability, Cronbach's $\alpha = 0.88$.

We also explored *state gustatory mental imagery* as a possible mediator of multisensory VR exposure, as effects may be contingent upon the extent to which an individual can mentally simulate the consumption of a food (Spence, 2011). *State gustatory mental imagery* was assessed in all conditions using three items (e.g., "To what extent were you able to imagine/picture yourself eating the chocolate") rated on a 100 mm VAS with endpoints "Not At All" to "Image as clear and vivid as real-life" (cf. Tiggemann & Kemps, 2005). The scale displayed good consistency, Cronbach's $\alpha = 0.95$.

2.5.3 Manipulation and debriefing checks. To assess whether our manipulation of hunger was successful, we asked participants to rate their *hunger* level at the beginning of test sessions on a 100 mm VAS (anchored from "Not at all" to "Very Much"). We similarly examined whether participants perceived the food stimulus (i.e., chocolate) as sufficiently rewarding by asking them to rate *liking* for chocolate on a 100 mm VAS (anchored from "Not At All" to "Very Much").

A debriefing check (Robinson et al., 2018) was performed at the end of test

sessions, to check if our implemented controls (i.e., cover story and distraction tasks) successfully diffused participants' awareness of study aims. The debriefing questionnaire probed study suspicions using a mixture of open-ended questions (e.g., "Did you hear about this study from other people, prior to participating"?) and one multiple-choice question. The latter required individuals to guess the real study aim from an array of 11 possible options.

2.5.4 Control variables. We examined one's *familiarity with VR technology* using a five-point Likert item (anchored from 1 = "I do not know of or recognize VR technology" to 5 = "I regularly use VR technology"; adapted from Tuorila *et al.*, 2001). Furthermore, demographic and anthropometric characteristics (i.e. *sex, age, self-reported height* and *weight*) were collected at the onset of testing. Self-reported height and weight were used to calculate participants' *BMI* (in kg/m²).

Finally, as exploratory individual-level correlates, we measured one's ability to form mental representations (i.e., trait mental imagery) arising from different sensory input: *Trait olfactory sensory imagery* (Cronbach's $\alpha = 0.76$) and *trait gustatory sensory imagery* (Cronbach's $\alpha = 0.68$). Both were assessed via respective (five-item) subscales of the Plymouth Sensory Imagery Questionnaire (Andrade *et al.*, 2014).

2.6 Data Analysis

We conducted all data analysis using R (version 4.4.1; R Core Team, 2024) and the *brms* package (Bürkner, 2017). Because we hypothesized parameter equality (H_{1D}), we adopted a Bayesian approach for our main analyses, as it facilitates assessing evidence for equivalence (i.e., treatment differences equal to zero). For our Bayesian models (H_{1A} to H_2 and exploratory counterparts; **sections 3.2** to **3.6**), to align our inferences with the traditional 0.05 cutoff value, we decided *a priori* to declare estimates as credibly non-zero, or *credible*, if their 95% confidence interval excluded zero (for non-directional hypotheses) or if their posterior probability of direction (p^+ : proportion of posterior draws in hypothesized direction; Bayesian equivalent to 1-one-sided *p*-value; Marsman & Wagenmakers, 2016) exceeded 95%. For non-Bayesian analyses (e.g., manipulation checks; cf. **section 3.1**), we used the traditional 0.05 cutoff value for statistical significance. We discuss modelling details below where appropriate. Our analyses were pre-registered (URL: <u>osf.io/6hjax</u>) and blinded, meaning that the dataset was initially analysed with anonymized condition labels.

3. Results

3.1 Manipulation and debriefing checks

First, we tested whether mean *liking* and *hunger* scores were different from the scale midpoint (50) with respective one-sample z-tests. *Liking* was significantly higher than the midpoint (76 versus 50 mm, 95% CI = [72, 80]), Z = 11.85, p < .01, indicating that participants perceived the food stimulus (i.e., chocolate) as sufficiently rewarding. Conversely, *hunger* ratings were not significantly higher than neutral (54 versus 50 mm, 95% CI = [48, 59]), Z = 1.21, p = 0.20. We additionally examined whether *hunger* scores were significantly different from those of an earlier investigation that employed the same hunger manipulation (cf. van der Waal *et al.*, 2021). A one-sample z-test revealed that participants' *hunger* level was significantly lower than the previous study (51 versus 62.95 mm, 95% CI = [46, 56]), Z = -4.45, p < .01. Taken together, these results suggest that hunger state was not successfully manipulated. Consequently, we made exploratory adjustments for hunger in our main models (cf. **section 3.4**) to circumvent potential floor effects on our primary research aims.

Debriefing checks with a one-sample exact binomial test showed that 20 out of 70 individuals (i.e., 29%, 95% CI = [.18, .41]) guessed the study's aims correctly. Consequently, we re-ran our main confirmatory models excluding individuals who correctly guessed study aims. Findings from these hypothesis tests did not change initial conclusions, so we will not discuss them further.

3.2 Psychological and physiological FCRs within exposure modes (H_{1A})

To examine whether *craving* and *salivary volume* between food and non-food stimuli (i.e., psychological and physiological FCRs, respectively) were credibly different across exposure modes (H_{1A}), we formulated a Bayesian multilevel regression model for each outcome measure. In each model, we specified main and interaction effects of *Stimulus Type* and *Exposure Mode* as fixed effects, *Participant* as a random effect, and *Familiarity with VR technology*, *Sex*, *Age*, and *BMI* as (centred) covariates in the fixed part of the model.

Results yielded a main effect of *Stimulus Type* on psychological FCRs: *Craving* was systematically higher for food versus non-food in Unisensory VR (Mean Difference F-NF: 12.43 mm, 95% CI = [6.45, 18.33]), Multisensory VR (Mean Difference F-NF: 17.21 mm, 95% CI = [11.31, 23.20]), as well as Real-life conditions (Mean Difference F-NF: 29.80 mm, 95% CI = [23.81, 35.59]), all $p^+ = 100.00\%$ (**Figure 3**).



Figure 3. Means and 95%CIs of craving (left) and salivary volume (right) across the experimental conditions. RL = Real Life, MVR = Multisensory Virtual Reality; UVR = Unisensory Virtual Reality; F = Food; NF = Non-food.

While *salivary volume* was greater on average after interacting with food versus nonfood stimuli in all exposure modes (**Figure 3**), an interaction between *Stimulus Type* and *Exposure Mode* was detected as physiological FCRs were only credible in the Real-life setting (Mean Difference F-NF: 0.08 g, 95% CI = [0.03, 0.13]), p^+ = 99.90%. Therefore, partial evidence was found for H_{1A}.

3.3 Psychological and physiological FCRs between exposure modes (Δ FCRs; H_{1B}-H_{1D})

We then examined differences in psychological and physiological FCRs (Δ FCRs) between exposure modes (H_{1B} – H_{1D}) from the model described above. As H_{1D} required testing for *equivalence* (i.e., zero treatment difference), it merited a different approach: We treated FCRs whose difference was within 0.1 standard deviations of the respective outcome (SD_{craving} = 30.13, SD_{volume} = 0.39) from zero as equivalent to one another. Then we calculated the proportion of the respective posterior distribution within that interval (i.e., region of practical equivalence; ROPE) to ascertain confidence in the equivalence statement (Kruschke, 2011; Kruschke, 2018). Note that this procedure is similar to the frequentist concept of equivalence testing, a practical method for assessing evidence for an interval-null hypothesis.

First, we expected that psychological and physiological FCRs would be stronger in Multisensory VR relative to Unisensory VR conditions (H_{1B}). As displayed in **Figure 4A** and **4B**, while differences were in the expected direction, Multisensory VR exposure did not trigger systematically higher food-specific *craving* (Mean Difference _{MVR-UVR}: 4.79 *mm*, 95% CI = [-3.52, 13.17]), $p^+ = 86.92\%$, nor greater *salivary volume* towards food versus non-food cues (Mean Difference _{MVR-UVR}: 0.02 g, 95% CI = [-0.05, 0.09]), $p^+ = 73.83\%$, compared to Unisensory VR settings. Thus, H_{1B} was not supported. That said, we also did not find evidence for equivalence (i.e., *zero* difference) in craving profiles between Multisensory VR and Unisensory VR exposures (**Figure 4B**).

Moreover, we anticipated that psychological and physiological FCRs would be greater in Real-life relative to Unisensory VR exposure (H_{1C}). Figure 4B shows that H_{1C}



Figure 4. A) Psychological (craving) and physiological (salivary volume) responses to food versus non-food cues (i.e., FCRs) across exposure modes. RL: Real Life. UVR: Unisensory VR. MVR: Multisensory VR. Points and intervals are posterior means and 95% CIs. Filled points denote credible (non-zero) differences between food and non-food stimuli. B) Differences in psychological and physiological FCRs (i.e., Δ FCRs) between different exposure modes. Dashed lines indicate limits of the region of practical equivalence (ROPE) to zero. ROPE is the percentage of the posterior distribution in the region of practical equivalence to zero, and thus assesses evidence for the null of no difference (i.e., *equivalence* between conditions). Points and intervals are posterior means and 95% CIs. Filled points denote credible (non-zero) differences between conditions.

was supported: In Real-life, the difference in *craving* between food versus non-food stimuli was indeed bigger compared to the Unisensory VR condition (Mean Difference _{RL-UVR}: 17.38 *mm*, 95% CI = [8.87, 25.67]), p^+ = 100.00%. Likewise, food-induced *salivary volume* was greater in Real-life than in Unisensory VR settings (Mean Difference _{RL-UVR}: 0.06 *g*, 95% CI = [-0.01, 0.13]), p^+ = 96.16%.

Finally, we assessed whether psychological and physiological FCRs were similar across Multisensory VR and Real-life conditions. Contrary to hypothesized (H_{1D}), Multisensory VR exposure elicited systematically weaker food-specific *craving* than in Reallife (Mean Difference _{RL-MVR}: 12.59 *mm*, 95% CI = [4.28, 20.95]), p^+ = 99.80%. On the other hand, and congruent with expectations, *salivary volume* towards food (versus non-food stimuli) did **not** credibly differ between Multisensory VR and Real-life settings (Mean Difference _{RL-MVR}: 0.04 *g*, 95% CI = [-0.03, 0.11]), p^+ = 87.79% – although we could not conclude equivalence (i.e., *zero* difference) in salivary profiles either (**Figure 4B**).

3.4. Exploratory corrections for hunger state

Given that earlier checks indicated an unsuccessful hunger manipulation, we explored

whether correcting for participants' hunger state would better distinguish psychological and physiological FCRs between exposure modes. Thus, we adjusted for hunger state by including a main effect of (grand mean-centered) *Hunger* and its interactions with *Stimulus Type* and *Exposure Mode* as predictors in otherwise identical models (cf. section 3.2).

In **Table 1**, we show differences in FCRs between exposure modes at one standard deviation (i.e., 24 units) above (+1 SD) and below (-1 SD) mean hunger. Results confirmed weaker food-specific *craving* in both virtual conditions relative to Real-life – even at matched levels of hunger. Similarly, a lower salivary response to food (versus non-food) in Unisensory VR compared to Real-life settings was replicated, as well as the previous finding that salivary profiles did not credibly differ between Multisensory VR and Real-life exposure. That said, correcting for hunger revealed an additional difference in physiological FCRs between virtual exposure modes in line with H_{1B} : Food-induced *salivary volume* was 0.11 units lower (95% CI = [-0.20, -0.01]) in the Unisensory VR compared to the Multisensory VR setting, even when individuals reported higher (+1 SD) hunger levels in the former condition. Interestingly, within Unisensory VR exposure, higher (+1 SD) hunger levels were associated with weaker salivary responses (B = -0.13, 95% CI = [-0.22, -0.03]; **Table 1**).

FCR Outcome Contrast		Estimate and 95% CI (<i>p</i> ⁺)	
Craving	UVR (hunger.1 SD) - RL (hunger.1 SD)	-14.16 [-25.45, -2.47] (99.21%)	
Craving	MVR (hunger-1 SD) - RL (hunger-1 SD)	-10.70 [-22.17, 0.69] (96.61%)	
Craving	MVR (hunger.1 SD) - UVR (hunger.1 SD)	3.46 [-8.30, 14.85] (72.07%)	
Craving	RL (hunger+1 sp) - RL (hunger-1 sp)	6.85 [-4.50, 18.34] (88.05%)	
Craving	RL (hunger _{+1 SD}) - UVR (hunger _{-1 SD})	21.01 [9.62, 32.25] (99.99%)	
Craving	RL (hunger _{+1 SD}) - MVR (hunger _{-1 SD})	17.55 [6.14, 29.11] (99.81%)	
Craving	UVR (hunger _{+1 SD}) - RL (hunger _{-1 SD})	-13.59 [-25.25, -2.02] (98.78%)	
Craving	UVR (hunger _{+1 SD}) - UVR (hunger _{-1 SD})	0.57 [-11.03, 12.15] (53.31%)	
Craving	UVR (hunger _{+1 SD}) - MVR (hunger _{-1 SD})	-2.90 [-14.52, 8.88] (69.19%)	

Table 1. Psychological and physiological FCRs between exposure modes (Δ FCRs) at specified levels of hunger (i.e., 1 SD above and below mean hunger).

Craving	UVR (hunger+1 sp) - RL (hunger+1 sp)	-20.44 [-32.08, -8.83] (99.97%)		
Craving	MVR (hunger+1 SD) - RL (hunger-1 SD)	-7.37 [-18.72, 4.09] (89.44%)		
Craving	MVR (hunger+1 sp) - UVR (hunger-1 sp)	6.79 [-4.89, 18.26] (87.69%)		
Craving	MVR (hunger+1 SD) - MVR (hunger-1 SD)	3.33 [-7.95, 14.74] (71.42%)		
Craving	MVR (hunger+1 sp) - RL (hunger+1 sp)	-14.22 [-25.68, -2.74] (99.34%)		
Craving	MVR (hunger _{+1 SD}) - UVR (hunger _{+1 SD})	6.22 [-5.54, 17.91] (84.85%)		
Salivary Volume	UVR (hunger-1 SD) - RL (hunger-1 SD)	0.00 [-0.09, 0.10] (53.30%)		
Salivary Volume	MVR (hunger _{-1 SD}) - RL (hunger _{-1 SD})	-0.02 [-0.11, 0.08] (64.14%)		
Salivary Volume	MVR (hunger-1 SD) - UVR (hunger-1 SD)	-0.02 [-0.12, 0.07] (67.08%)		
Salivary Volume	RL (hunger _{+1 SD}) - RL (hunger _{-1 SD})	0.01 [-0.09, 0.10] (56.07%)		
Salivary Volume	RL (hunger+1 SD) - UVR (hunger-1 SD)	0.00 [-0.09, 0.10] (52.52%)		
Salivary Volume	RL (hunger _{+1 SD}) - MVR (hunger _{-1 SD})	0.02 [-0.07, 0.12] (69.92%)		
Salivary Volume	UVR (hunger+1 sp) - RL (hunger-1 sp)	-0.13 [-0.22, -0.03] (99.52%)		
Salivary Volume	UVR (hunger _{+1 SD}) - UVR (hunger _{-1 SD})	-0.13 [-0.22, -0.03] (99.45%)		
Salivary Volume	UVR (hunger _{+1 SD}) - MVR (hunger _{-1 SD})	-0.11 [-0.20, -0.01] (98.60%)		
Salivary Volume	UVR (hunger _{+1 SD}) - RL (hunger _{+1 SD})	-0.13 [-0.23, -0.04] (99.60%)		
Salivary Volume	MVR (hunger+1 SD) - RL (hunger-1 SD)	-0.06 [-0.16, 0.04] (88.88%)		
Salivary Volume	MVR (hunger _{+1 SD}) - UVR (hunger _{-1 SD})	-0.06 [-0.16, 0.03] (90.50%)		
Salivary Volume	MVR (hunger+1 SD) - MVR (hunger-1 SD)	-0.04 [-0.14, 0.05] (80.83%)		
Salivary Volume	MVR (hunger _{+1 SD}) - RL (hunger _{+1 SD})	-0.07 [-0.16, 0.03] (91.71%)		
Salivary Volume	MVR (hunger+1 SD) - UVR (hunger+1 SD)	0.07 [-0.03, 0.16] (90.71%)		

3.5 Indirect mediation of Multisensory VR effects by presence (H₂)

In light of (directional) improvements in food-induced *craving* and *salivary volume* afforded by Multisensory VR over Unisensory VR exposure, we further examined potential psychological processes that might mediate these differences (H₂). To this end, we first calculated mean FCRs for each virtual exposure mode (across *Stimulus Type*), because *presence* (the hypothesized mediator) was measured once per virtual condition. We then specified a multivariate regression model predicting *Presence* from virtual *Exposure Mode* (i.e., Unisensory VR versus Multisensory VR; E -> P), and one predicting FCRs (i.e., *craving* or *salivary volume*) from *Presence* and all covariates as specified in the previous model (P -> FCRs; **Table 2**). We specified both paths (i.e., E ->P and P -> FCR) with by-person random intercepts to account for repeated measures over individuals. We then quantified whether *presence* mediates FCR effects of the Multisensory VR condition by multiplying the virtual *Exposure Mode* to *Presence* path coefficient with the *Presence* to FCR path coefficient (i.e., E->P * P->FCR; **Table 2**).

The mediation model indicated that shifting from the Unisensory VR to Multisensory VR condition increased feelings of *presence* (cf. **Table 2** and **Figure 5A**). However, *presence* ratings did not further predict either *craving* or *salivary volume* (**Figure 5B and 5C**). In other words, while Multisensory VR exposure did enhance experienced *presence* in the virtual environment, the latter did not (indirectly) account for effects on either psychological or physiological FCRs. We therefore did not find support for H₂.

3.6 Exploratory mediation by state gustatory mental imagery

We explored whether (subtle) differences in FCR profiles between virtual conditions could instead be explained by variations in the extent to which one could mentally simulate the consumption of a food (i.e., state gustatory mental imagery). To this end, we formulated an identical mediation model as above (H₂) but with *state gustatory mental imagery* as the mediator.

Results showed that Multisensory VR exposure significantly enhanced perceptions of *state gustatory mental imagery* compared to the Unisensory VR counterpart (cf. **Table 2** and **Figure 5D**). In turn, increased *state gustatory mental imagery* predicted greater *craving* ratings (**Figure 5E**), but did not correlate with *salivary volume* (**Figure 5F**). However, *state gustatory mental imagery* ratings did correlate positively with individuals' (baseline) abilities to mentally simulate tasting (i.e., *trait gustatory mental imagery*; r (68) = 0.38, t = 3.39, p < 0.01), as well as smelling products (i.e., *olfactory mental imagery*; r = 0.35 (68), t = 3.12, p < 0.01

0.01).

Table 2. Mediation model results for psychological and physiological FCRs across virtual exposure modes.

FCR Outcome	Path	Mean	SD	95% CI	p^+					
Model 1: Indirect mediation via presence (H ₂)										
Craving ¹	E->P	0.28	0.05	[0.19, 0.38]	100.00%					
Craving	P->FCR	8.13	6.14	[-4.01, 20.27]	90.87%					
Craving	E->P * P->FCR	2.30	1.81	[-1.12, 6.06]	90.87%					
Salivary Volume	P->FCR	-0.05	0.06	[-0.17, 0.06]	82.98%					
Salivary Volume	E->P * P->FCR	-0.02	0.02	[-0.05, 0.02]	82.98%					
Model 2: Indirect mediation via state gustatory mental imagery (Exploratory)										
Craving ¹	E->I	18.58	2.46	[13.72, 23.41]	100.00%					
Craving	I->FCR	0.61	0.05	[0.52, 0.70]	100.00%					
Craving	E->I * I->FCR	11.35	1.74	[8.08, 14.88]	100.00%					
Salivary Volume	I->FCR	0.00	0.00	[0.00, 0.00]	56.06%					
Salivary Volume	E->I * I->FCR	0.00	0.01	[-0.02, 0.02]	56.06%					

¹Path is shown only once because conceptual mediators (i.e., *presence* and *state gustatory mental imagery*) were measured per virtual condition and thus aggregated across FCR outcomes. E: Exposure Mode (Unisensory VR versus Multisensory VR), P: Presence ratings (five-point Likert scale ranging from 1 to 5), I: State gustatory mental imagery ratings (100 mm VAS).



Figure 5. Top row: Results of the mediation model assessing whether *presence* accounted for effects of Multisensory VR exposure on psychological (i.e., craving) and physiological FCRs (i.e., salivary volume). A) Conditional posterior means and 95%CIs of *presence* in Unisensory (UVR) and Multisensory (MVR) VR exposure modes. B) Model's regression line and 95%CI of *craving* on *presence*. C) Model's regression line and 95%CI of *salivary volume* on *presence*. Bottom row: Results of the mediation model that explored whether *state gustatory mental imagery* explained effects of Multisensory VR (MVR) exposure on *craving* and *salivary volume*. D-F) As A-C except with *state gustatory mental imagery* as mediator.

4. Discussion

The present lab-based study systematically investigated whether an immersive multisensory VR food environment – with the added presence of olfactory cues – could validly model individuals' fundamental (psychological and physiological) responses to food, thereby bridging the current gap between existing "vision-only" unisensory VR environments and real-life settings. In sum, we found that food cues induced stronger craving than non-food cues across all exposure modes, while this was only true for salivary volume in the (physical) Real-life condition (H_{1A}). Additionally, Unisensory VR exposure consistently elicited weaker psychological and physiological FCRs compared to Real-life (H_{1C}), whereas Multisensory

VR exposure led to a directional (non-equivalent) improvement over Unisensory VR settings in food-specific craving and higher salivary food responses after adjusting for hunger (H_{1B}). We could not conclude equivalence between Multisensory VR and Real-life conditions on either FCR outcome, though the latter did not systematically differ from one another on the basis of salivary responses either (H_{1D}). Finally, although Multisensory VR exposure increased feelings of presence, this increase could not account for the differences in FCRs that emerged between virtual conditions. Instead, subtle differences in craving profiles between virtual conditions were mediated by enhanced gustatory mental imagery in the Multisensory VR setting.

The psychological FCR, craving, showed strong and robust effects of the food versus non-food manipulation, with stronger craving for food cues across all exposure modes. This main effect of stimulus type also remained significant after exploratory corrections for different levels of hunger state. As such, our results replicate the findings of van der Waal *et al.* (2021), who demonstrated that *within* an exposure mode, (visual) food cues reliably led to stronger reported craving in both virtual and real-life settings. We now expand on these results by revealing that (visual + olfactory) food cues also lead to higher craving than non-food cues for the multisensory VR condition, which was in line with our expectations.

Besides a main effect of stimulus type, we also observed differences in food-specific craving *between* different exposure modes in our study. We replicate another result by van der Waal *et al.* (2021), showing that while food-induced craving was credible in all exposure modes, its expression was consistently weaker in virtual settings relative to real-life. We anticipated that olfactory cues in the multisensory VR condition would boost one's desire to consume the food, because of previous findings that food odours alone can trigger one's appetite and particularly craving for the corresponding food, as well as foods with similar

taste profiles (i.e., "sensory-specific appetite"; Ramaekers *et al.*, 2014; Wolz *et al.*, 2017). However, in line with the meta-analysis of Boswell & Kober (2016), the addition of olfactory cues did not confer a clear benefit over using solely visual food cues for eliciting craving. One possibility for this null finding is that the odour was not intense enough to sufficiently induce participants' food consumption beliefs. Namely, the intensity of an odour indicates one's proximity to a food source (Jacobs, 2012), and a stronger intensity may have signalled a closer distance to the associated product (i.e., chocolate), thereby strengthening beliefs surrounding how easily attainable it is to subsequently consume (Spence, 2011). That said, the flow rate and volume of odor solution contained within *Sniff-O* were carefully piloted to be at a medium-to-high intensity odour level across individuals. We therefore do not think that odour exposure was too weak, in principle, to elicit an observable effect on food-specific craving.

An alternative explanation for the absence of notable differences between multisensory and unisensory VR settings in craving may be a lack of motivation – rather than ability or opportunity – to use olfactory information in one's cognitive processing of foods. Research has suggested that one's importance attached to the sense of smell (Croy *et al.*, 2009), general attendance to olfactory stimuli in the environment (Smeets *et al.*, 2008), and consequent application of olfactory information in decision making (Koller *et al.*, 2023) varies across individuals. Importantly, there is also evidence of cross-cultural variations such that those from industrialized English-speaking cultures (such as our participant sample) use the olfactory sense less frequently and readily compared to the visual sense (Majid, 2021; Majid *et al.*, 2018). Indeed, a dominance or "primacy" of the visual sense has been documented (among sighted-individuals) during moments of exploring a product, as well as upon food choice (Fenko *et al.*, 2010; Schifferstein *et al.*, 2013).

It was the case, though, that food-induced craving was directionally higher after

multisensory VR exposure, and equivalence also could not be established among virtual conditions, suggesting that the added odour was at least partially (albeit not systematically) successful at increasing craving perceptions. Craving is a marker for one's reactivity to food cues in the environment (Higgs, 2016; Nederkoorn, 2000), and has shown to be a moderate predictor of, and consequent therapeutic target for, an individual's dietary intake (Boswell & Kober, 2016). Taken together, our results therefore support the notion that exposure to food cues in VR can be successful at inducing ecologically-relevant psychological food responses such as craving, albeit to a smaller degree than as experienced in real-life.

For physiological food cue responses, our study showed increased salivary responses to real food versus non-food cues, thereby confirming the potency of (real) chocolate stimuli to robustly induce detectable salivary responses that serve to prepare the body for ingestion. Further, in line with our expectations and previous findings of van der Waal et al. (2021), we found that food-induced salivary responses were weaker in unisensory VR than in real-life The expected benefit of additional smell cues, which we anticipated to reduce this gap in salivation between real-life and virtual settings, because of the importance of olfaction in appetitive responses (Morquecho-Campos et al. 2019), was less evident. Though we could not establish that multisensory VR exposure led to equivalent food-induced salivation as in real-life, there was also no significant difference in salivary responses between these conditions, suggesting that there may be a directionally beneficial effect of adding olfactory cues. Notably, after adjusting for hunger levels-an important motivational factor in foodrelated behaviors and responses, as indicated by previous research (Rogers and Hill, 1989), and considered here due to the unsuccessful hunger manipulation-differences in salivary responses emerged between unisensory and multisensory VR conditions. This finding highlights hunger as a potential boundary condition that influences the effectiveness of

multisensory VR applications, suggesting that the added benefit of multisensory stimulation in VR may be contingent on an individual's hunger state.

Exploring possible mechanisms underpinning effects of multisensory VR exposure on FCRs, we demonstrate that state gustatory mental imagery mediates the subtle difference in craving responses between multisensory versus unisensory VR conditions. State mental imagery is the extent to which participants can imagine, in that specific trial, what eating the food would be like (e.g., 'To what extent were you able to imagine yourself eating the chocolate'). Our study thus supports prior work showing that mental imagery, and particularly individuals' ability to cognitively elaborate on the taste of a product, is a prerequisite for developing craving for a food (Croijmans & Wang, 2021; Higgs, 2016). Our findings further underscore that multisensory VR exposure does affect cognitive states, as it led participants to experience stronger state gustatory mental imagery. This in turn predicted higher levels of craving, which may facilitate later choice for the food, by shifting one's mindset towards obtaining the target of craving (Boswell & Kober, 2016; Higgs, 2016; Muñoz-Vilches *et al.*, 2020). For instance, enhancing mental simulations by manipulating sensory aspects in pictures was found to increase product liking and purchase intention in marketing (Krishna *et al.*, 2016).

We did not observe a similar mediation of state mental imagery on salivation levels, in contrast to previous research showing that self-reported mental imagery vividness influences salivary control (White, 1978), and that imagining a food odour - in combination with the image of a food - increases salivary responses (Krishna *et al.*, 2014). However, we did find that state (gustatory) mental imagery was associated with one's trait mental imagery, in the sense that the ability to experience imagery, in different sensory modalities, is a trait that varies across individuals (Zeman *et al.*, 2020). An individual's trait gustatory and olfactory imagery were both positively related to state mental imagery, stressing that baseline

imagery ability can be a relevant moderator of the effectiveness of VR applications in the food domain. Studies have revealed that people with higher (trait) mental imagery ability tend to imagine a product's scent spontaneously upon seeing an image (Sharma & Estes, 2024), need less elaborate descriptions of products than low imagers to experience the same level of craving (Croijmans & Wang, 2021), and report higher levels of craving after exposure to smell alone (Krishna, 2014). Our results are compatible with these earlier observations, and importantly, demonstrate that mental imagery strength (whether state or trait) can be intrinsic to effects of multisensory food cue exposure in a VR setting.

It is worth noting that mental imagery vividness has been positively linked to the sense of presence in VR environments (Iachini et al., 2018), yet our data show that presence was not correlated with any FCR outcomes. Presence captures the extent to which a user suspends (physical) reality and feels part of the (virtual) environment (Slater & Wilbur, 1997). As a central conceptual mediator of various VR applications, a greater sense of presence is widely anticipated to generate results that more accurately reflect measurements obtained in real physical settings. For example, liking ratings obtained in a multisensory VR coffeehouse was more predictive of future coffee liking than ratings obtained from traditional sensory booths in a laboratory (Bangcuyo et al., 2015). The majority of work in VR thus focuses on how to enhance presence, by increasing technological immersion (e.g., through delivering a multisensory experience), or by creating opportunities for natural interaction (Bowman & McMahan, 2007). While it is well-documented that higher technological immersion enhances feelings of presence (Cummings & Bailenson, 2014), establishing a connection between higher levels of presence and a plethora of (cognitive) outcomes such as learning has proven challenging (Makransky et al., 2017). Our results therefore reflect the current state of evidence in that we indeed found systematic improvements in presence with additional sensory (olfactory) information, but the subsequent impact of presence on FCRs

did not manifest. This indicates that regardless of how well constructed a virtual environment is, there is still a need for individuals to accept the premise of the virtual experience for it to be sufficiently effective. We require more empirical studies on not only the role of presence as a mediator, but also a refined theoretical framework for explaining and predicting the effects of higher levels of presence (Barranco Merino *et al.*, 2023).

This study is not without its limitations. First, a minority of participants (N = 11; 16%) did not adhere to pre-test fasting instructions, which was also echoed in failed manipulation checks for hunger. This may have dampened an individual's overall reactivity to food (versus non-food) cues and unintendedly introduced floor effects on craving and salivation. Indeed, if we did not perform exploratory corrections for hunger ratings, we would not have uncovered the further difference in food-induced salivary responses between unisensory VR and multisensory VR exposure. That said, it is worth noting that while hunger ratings were not significantly higher than the neutral (midpoint) value, liking ratings for the food stimulus (i.e., chocolate) were. Collectively, we can therefore assume that individuals perceived food cues to be sufficiently rewarding (Berridge, 2009). Second, debriefing checks revealed that 20 (29%) participants correctly guessed the purpose of the research, which could have triggered a tendency for these respondents to report socially desirable craving ratings (Orne, 2017). However, awareness of study aims was likely less problematic for our physiological (salivation) outcome, whose activity is much less subject to conscious control (Dawes, 1996). Finally, a few incongruencies were present between virtual and real-life environments that could have hindered accurate comparisons between exposure modes. For the real-life condition, we replicated the promotional product set-up in a controlled lab setting, as we prioritized internal validity for this "proof-of-concept" study. Though an actual supermarket setting would have been favourable for enhancing correspondences between exposure modes, it would have required extra resources (e.g., a larger participant sample) to

account for additional sources of "noise" in the data (e.g., competing sensory input from nearby products and people). Similarly, in virtual settings we used a non-parametric version of the *Sniff-O* olfactometer that administered a constant intensity of an odour – irrespective of the distance or movement trajectory of the associated virtual object. The latter may have likewise limited comparisons with real-life circumstances, in which odours diffuse in space and create intensity gradients that vary in proximity to their source (Jacobs, 2012).

Lastly, there are a few unresolved questions that require further investigation. As multisensory VR environments become a frontier topic in the development of next generation immersive technologies (Melo et al., 2022), it would be worthwhile to explore its promises for domains that traditionally proved difficult for virtual experiences, such as (public) health research. For instance, future studies could investigate the utility of such a multisensory VR supermarket for measuring other physiological (e.g., salivary enzyme activity; Morquecho-Campos et al., 2019) and behavioural markers (e.g., food spatial memory and grocery purchases; de Vries et al., 2021) of dietary relevance - especially if such a VR infrastructure can incorporate additional sources of data from emerging components like heart rate monitors and eye-tracking (Halbig & Latoschik, 2021). If the latter were achieved, we could use the multisensory VR set-up to model the impact of potential health interventions, such as simulating how structural (policy) changes to the physical food environment (e.g., nudging) can impact dietary choices (Blom et al., 2021; Larson & Story, 2009), across different socioeconomic groups (Mizdrak et al., 2017). The merits of such a set-up for non-food purposes can likewise be assessed, such as for cue exposure addiction therapy and smell training in clinical populations (Hone-Blanchet et al., 2014; Hwang et al., 2023).

In conclusion, this pre-registered lab experiment highlights that an immersive multisensory VR food environment with olfactory cues can credibly model psychological (craving) responses to foods, albeit to a weaker degree than experienced in real-life. The

added value of such a technology over unisensory (vision-only) VR infrastructures may lie in enhancing conceptual mediators (i.e., presence and consumption mental imagery) and in approaching physiological (salivary) responses to foods as in real-life. However, important boundary conditions (e.g., hunger state, trait mental imagery) should be taken into account to maximize these benefits.

Acknowledgements

This project was co-funded by the Digital Sciences for Society growth project grant program of Tilburg University (grant number: DSFS 2023011). The funding body was not involved in the research itself, or in the writing and submission of this article. Many thanks to *CyNexo srl* for providing the olfactory "Sniff-O" device to use in the multisensory VR environment. We would also like to thank Emma van Aken for help with data collection. **Declaration of interests: none.**

Author Contributions

RdV, NvdL, TvL, MV, and SB conceptualized the theoretical framework and study design, as well as acquired funding for the research. RdV and NvdL developed study materials. KV assisted with data collection and data curation, under the supervision of RdV. MV analysed and visualized the data. All authors interpreted study data, contributed to writing and editing of the manuscript, and approved the final article.

References

- Albayay, J., Castiello, U., & Parma, V. (2022). The effect of odour valence and odour detection threshold on the withholding and cancellation of reach-to-press responses. *Chemosensory Perception*, 1-14.
- Andonova, V., Reinoso-Carvalho, F., Jimenez Ramirez, M. A., & Carrasquilla, D. (2023).
 Does multisensory stimulation with virtual reality (VR) and smell improve learning?
 An educational experience in recall and creativity. *Frontiers in psychology, 14,* 1176697. https://doi.org/10.3389/fpsyg.2023.1176697
- Andrade, J., May, J., Deeprose, C., Baugh, S. J., & Ganis, G. (2014). Assessing vividness of mental imagery: The Plymouth sensory imagery questionnaire. *British Journal of Psychology*, 105(4), 547-563.
- Archer, N. S., Bluff, A., Eddy, A., Nikhil, C. K., Hazell, N., Frank, D., & Johnston, A.
 (2022). Odour enhances the sense of presence in a virtual reality environment. *PloS One*, 17(3), e0265039. https://doi.org/10.1371/journal.pone.0265039
- Bangcuyo, R. G., Smith, K. J., Zumach, J. L., Pierce, A. M., Guttman, G. A., & Simons, C. T. (2015). The use of immersive technologies to improve consumer testing: The role of ecological validity, context and engagement in evaluating coffee. *Food Quality and Preference*, 41, 84-95.
- Barranco Merino, R., Higuera-Trujillo, J. L., & Llinares Millán, C. (2023). The Use of Sense of Presence in Studies on Human Behavior in Virtual Environments: A Systematic Review. *Applied Sciences*, 13(24), 13095. https://doi.org/10.3390/app132413095
- Baus, O., Bouchard, S. (2017) Exposure to an unpleasant odour increases the sense of Presence in virtual reality. *Virtual Reality*, 21, 59-74.

Baus, O., Bouchard, S., Nolet, K. (2018). Exposure to a pleasant odour may increase the

sense of reality, but not the sense of presence or realism. *Behavior & Information Technology*, *38*, 1369-1378.

- Baus, O., Bouchard, S., Nolet, K., & Berthiaume, M. (2022). In a dirty virtual room: exposure to an unpleasant odor increases the senses of presence, reality, and realism. *Cogent Psychology*, 9(1), 2115690.
- Berridge, K. C. (2009). 'Liking'and 'wanting'food rewards: brain substrates and roles in eating disorders. *Physiology & Behavior*, 97(5), 537-550.
- Blom, S. S. A. H., Gillebaart, M., De Boer, F., van der Laan, N., & De Ridder, D. T. D.
 (2021). Under pressure: Nudging increases healthy food choice in a virtual reality supermarket, irrespective of system 1 reasoning. *Appetite*, 160, Article 105116. https://doi.org/10.1016/j.appet.2021.105116
- Boswell, R. G., & Kober, H. (2016). Food cue reactivity and craving predict eating and weight gain: a meta-analytic review. *Obesity reviews*, *17*(2), 159-177.
- Bowman, D. A., & McMahan, R. P. (2007). Virtual reality: How much immersion is enough? *Computer*, 40(7), 36–43.
- Brengman, M., Willems, K., & De Gauquier, L. (2022). Customer Engagement in Multi-Sensory Virtual Reality Advertising: The Effect of Sound and Scent Congruence. *Frontiers in Psychology*, 13, 747456. https://doi.org/10.3389/fpsyg.2022.747456
- Brunstrom, J. M., Yates, H. M., & Witcomb, G. L. (2004). Dietary restraint and heightened reactivity to food. *Physiology & Behavior*, 81(1), 85-90.
- Burgess, C. R., Ramesh, R. N., Sugden, A. U., Levandowski, K. M., Minnig, M. A., Fenselau, H., ... & Andermann, M. L. (2016). Hunger-dependent enhancement of food cue responses in mouse postrhinal cortex and lateral amygdala. *Neuron*, 91(5), 1154-1169.

- Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. Journal of Statistical Software, 80, 1–28. https://doi.org/10.18637/jss.v080.i01
- Cheah, C. S. L., Barman, S., Vu, K. T. T., Jung, S. E., Mandalapu, V., Masterson, T. D., . . . Gong, J. (2020). Validation of a Virtual Reality Buffet environment to assess food selection processes among emerging adults. *Appetite*, 153, 104741. https://doi.org/10.1016/j.appet.2020.104741
- Croijmans, I., & Wang, Q. J. (2022). Do you want a description with that wine? The role of wine mental imagery in consumer's desire to drink using the revised Vividness of Wine Imagery Questionnaire (VWIQ-II). *Journal of Sensory Studies*, *37*(1), e12712.
- Croy, I., Buschhüter, D., Seo, H. S., Negoias, S., & Hummel, T. (2010). Individual significance of olfaction: development of a questionnaire. *European Archives of Oto-Rhino-Laryngology*, 267, 67-71.
- Cummings, J. J., & Bailenson, J. N. (2014). How Immersive Is Enough?: A Meta-Analysis of the Effect of Immersive Technology on User Presence. *Media Psychology*, 19(2), 272– 309. https://doi.org/10.1080/15213269.2015.1015740
- Dawes, C. (1975). Circadian rhythms in the flow rate and composition of unstimulated and stimulated human submandibular saliva. *Journal of Physiology*, 244, 535-548.
- Dawes, C. (1987). Physiological factors affecting salivary flow rate, oral sugar clearance, and the sensation of dry mouth in man. *Journal of Dental Research*, 66, 648-653.
- Dawes, C. (1996). Factors influencing salivary flow rate and composition. *Saliva and Oral Health*, *2*, 27-41.
- Fenko, A., Schifferstein, H. N., & Hekkert, P. (2010). Shifts in sensory dominance between various stages of user–product interactions. *Applied ergonomics*, *41*(1), 34-40.

Gallace, A., Ngo, M. K., Sulaitis, J., and Spence, C. (2012). "Multisensory presence in virtual

reality: possibilities & limitations," in *Multiple Sensorial Media Advances andApplications: New Developments* in MulSeMedia, eds G. Ghinea, F. Andres, and S.R. Gulliver (Hershey, PA: IGI Global), 1–38.

- Goncalves, M., Vasconcelos-Raposo, M., Bessa, M. (2020). Impact of Different Sensory Stimuli on Presence in Credible Virtual Environments. *IEEE Transactions on Visualization and Computer Graphics*, *26*(11), 3231-3240.
- Grassini, S., & Laumann, K. (2020). Questionnaire measures and physiological correlates of presence: A systematic review. *Frontiers in Psychology*, *11*, 349.
- Halbig, A., & Latoschik, M. E. (2021). A Systematic Review of Physiological Measurements, Factors, Methods, and Applications in Virtual Reality. *Frontiers in Virtual Reality*, 2. https://doi.org/10.3389/frvir.2021.694567
- van Herpen, E., Van den Broek, E., Van Trijp, H. C. M. Yu, T. (2016) Can a virtual supermarket bring realism into the lab? Comparing shopping behavior using virtual and pictorial store representations to behavior in a physical store. *Appetite*, 107, 196-207.
- Higgs, S. (2016). Cognitive processing of food rewards. Appetite, 104, 10-17.
- Hill, A. J. (2007). The psychology of food craving: Symposium on 'Molecular mechanisms and psychology of food intake'. *Proceedings of the Nutrition Society*, *66*(2), 277-285.
- Hone-Blanchet, A., Wensing, T., & Fecteau, S. (2014). The use of virtual reality in craving assessment and cue-exposure therapy in substance use disorders. *Frontiers in human neuroscience*, 8, 844.
- Hummel, T., Kobal, G., Gudziol, H., & Mackay-Sim, A. J. E. A. (2007). Normative data for the "Sniffin'Sticks" including tests of odour identification, odour discrimination, and olfactory thresholds: an upgrade based on a group of more than 3,000 subjects. *European Archives of Oto-Rhino-Laryngology*, 264, 237-243.

- Hwang, S. H., Kim, S. W., Basurrah, M. A., & Kim, D. H. (2023). The efficacy of olfactory training as a treatment for olfactory disorders caused by coronavirus disease-2019: a systematic review and meta-analysis. *American Journal of Rhinology & Allergy*, 37(4), 495-501.
- Iachini, T., Maffei, L., Masullo, M., Senese, V. P., Rapuano, M., Pascale, A., ... & Ruggiero, G. (2019). The experience of virtual reality: are individual differences in mental imagery associated with sense of presence?. *Cognitive processing*, 20(3), 291-298.
- Jacobs, L. F. (2012). From chemotaxis to the cognitive map: the function of olfaction. *Proceedings of the National Academy of Sciences*, 109(supplement_1), 10693-10700.
- Kanoski, S. E., & Boutelle, K. N. (2022). Food cue reactivity: Neurobiological and behavioral underpinnings. *Reviews in Endocrine and Metabolic Disorders*, 23(4), 683-696.
- Kemps, E., & Tiggemann, M. (2009). Attentional bias for craving-related (chocolate) food cues. *Experimental and Clinical Psychopharmacology*, 17(6), 425–433. https://doi.org/10.1037/a0017796

Koller, M., Salzberger, T., Floh, A., Zauner, A., Sääksjärvi, M., & Schifferstein, H. N. (2023).

Measuring individual differences in active smelling to evaluate products–The ENFAS-Instrument. *Food Quality and Preference*, *110*, 104925.

- Krishna, A., Morrin, M., & Sayin, E. (2014). Smellizing cookies and salivating: A focus on olfactory imagery. *Journal of Consumer Research*, *41*(1), 18-34.
- Krishna, A., Cian, L., & Sokolova, T. (2016). The power of sensory marketing in advertising. *Current Opinion in Psychology*, *10*, 142-147.
- Kruschke, J. K. (2011). Bayesian assessment of null values via parameter estimation and model comparison. *Perspectives on Psychological Science*, *6*(3), 299-312.

- Kruschke, J. K. (2018). Rejecting or accepting parameter values in Bayesian estimation. Advances in methods and practices in psychological science, 1(2), 270-280.
- van der Laan, L. N., Papies, E. K., Ly, A., & Smeets, P. A. M. (2022). Examining the neural correlates of goal priming with the NeuroShop, a novel virtual reality fMRI paradigm. *Appetite*, 170, 105901.
- Larson, N., & Story, M. (2009). A review of environmental influences on food choices. *Annals of Behavioral Medicine*, 38(suppl 1), s56-s73.
- Loeber, S., Grosshans, M., Herpertz, S., Kiefer, F., & Herpertz, S. C. (2013). Hunger modulates behavioral disinhibition and attention allocation to food-associated cues in normal-weight controls. *Appetite*, *71*, 32–39.

https://doi.org/10.1016/j.appet.2013.07.008

- Lombart, M. & Ditton, T. (1997) At the heart of it all: The concept of presence. *Journal of Computer-mediated Communication*, 3 (2).
- Long, J. W., Pritschet, S. J., Keller, K. L., Cheah, C. S. L., Boot, L., Klippel, A., . . . Masterson, T. D. (2023). Portion size affects food selection in an immersive virtual reality buffet and is related to measured intake in laboratory meals varying in portion size. *Appetite*, 191, 107052. https://doi.org/10.1016/j.appet.2023.107052
- Lundström, J. N., Gordon, A. R., Alden, E. C., Boesveldt, S., & Albrecht, J. (2010). Methods for building an inexpensive computer-controlled olfactometer for temporally-precise experiments. *International Journal of Psychophysiology*, 78(2), 179-189.
- Majid, A. (2021). Human olfaction at the intersection of language, culture, and biology. *Trends in Cognitive Sciences*, *25*(2), 111-123.
- Majid, A., Roberts, S. G., Cilissen, L., Emmorey, K., Nicodemus, B., O'grady, L., ... & Levinson, S. C. (2018). Differential coding of perception in the world's languages. *Proceedings of the National Academy of Sciences*, *115*(45), 11369-11376.

- Makransky, G., Terkildsen, T. S., & Mayer, R. E. (2017). Adding immersive virtual reality to a science lab simulation causes more presence but less learning. *Learning and Instruction*, 60, 225–236. https://doi.org/10.1016/j.learninstruc.2017.12.007
- Marsman, M., & Wagenmakers, E.-J. (2016). Three Insights from a Bayesian Interpretation of the One-Sided P Value. *Educational and Psychological Measurement*. <u>https://doi.org/10.1177/0013164416669201</u>
- Melo, M., Gonçalves, G., Monteiro, P., Coelho, H., Vasconcelos-Raposo, J., & Bessa, M. (2022). Do Multisensory Stimuli Benefit the Virtual Reality Experience? A Systematic Review. *IEEE Transactions on Visualization and Computer Graphics*, 28(2), 1428–1442. https://doi.org/10.1109/TVCG.2020.3010088
- Mizdrak, A., Waterlander, W. E., Rayner, M., & Scarborough, P. (2017). Using a UK virtual supermarket to examine purchasing behavior across different income groups in the United Kingdom: development and feasibility study. *Journal of medical Internet research*, 19(10), e7982.
- Morquecho-Campos, P., Bikker, F. J., Nazmi, K., de Graaf, K., Laine, M. L., & Boesveldt, S. (2019). Impact of food odours signaling specific taste qualities and macronutrient content on saliva secretion and composition. *Appetite*, *143*, 104399.
- Morquecho-Campos, P., Hellmich, I. M., Zwart, E., de Graaf, K., & Boesveldt, S. (2022).
 Does odour priming influence snack choice? An eye-tracking study to understand food choice processes. *Appetite*, 168, 105772.
 https://doi.org/10.1016/j.appet.2021.105772
- Motoki, K., Togawa, T. (2022) Multiple senses influencing healthy food preference. *Current Opinion in Behavioral Sciences, 48*, 101223.

Muñoz-Vilches, N. C., van Trijp, H. C., & Piqueras-Fiszman, B. (2020). Tell me what you

imagine and I will tell you what you want: The effects of mental simulation on desire and food choice. *Food Quality and Preference*, *83*, 103892.

Munyan III, B. G., Neer, S. M., Beidel, D. C., & Jentsch, F. (2016). Olfactory stimuli increase

presence in virtual environments. PloS one, 11(6), e0157568.

- Navazesh, M. (1993). Methods for collecting saliva. *Annals of the New York Academy of Sciences*, 694, 72–77. <u>https://doi.org/10.1111/j.1749-6632.1993.tb18343.x.</u>
- Nederkoorn, C., Smulders, F. T. Y., Jansen, A. (2000) Cephalic phase responses, craving and food intake in normal subjects. *Appetite*, *35*, 45-55.
- Niedenthal, S., Fredborg, W., Lundén, P., Ehrndal, M., & Olofsson, J. K. (2023). A graspable olfactory display for virtual reality. *International Journal of Human-Computer Studies*, 169, 102928. https://doi.org/10.1016/j.ijhcs.2022.102928
- Olofsson, J. K., Niedenthal, S., Ehrndal, M., Zakrzewska, M., Wartel, A., & Larsson, M.
 (2017). Beyond smell-o-vision: Possibilities for smell-based digital media. *Simulation*& *Gaming*, 48(4), 455–479. <u>https://doi.org/10.1177/1046878117702184</u>
- Orne, M. T. (2017). On the social psychology of the psychological experiment: With particular reference to demand characteristics and their implications. In *Sociological methods* (pp. 279-299). Routledge.
- Persky, S., Dolwick, A. P. (2020). Olfactory Perception and Presence in a Virtual Reality Food Environment. *Frontiers in Virtual Reality*, 1, 571812. https://doi.org/10.3389/frvir.2020.571812
- Proserpio, C., de Graaf, C., Laureati, M., Pagliarini, E., & Boesveldt, S. (2017). Impact of ambient odours on food intake, saliva production and appetite ratings. *Physiology & Behavior*, 174, 35-41.

R Core Team. (2024). R: A Language and Environment for Statistical Computing. Version

4.4.1 (Version 4.4.1) [Computer software]. *R Foundation for Statistical Computing*. https://www.R-project.org/

- Ramaekers, M. G., Boesveldt, S., Lakemond, C. M., van Boekel, M. A., & Luning, P. A. (2014). Odors: appetizing or satiating? Development of appetite during odor exposure over time. *International journal of obesity* (2005), 38(5), 650–656. https://doi.org/10.1038/ijo.2013.143
- Robinson, E., Bevelander, K. E., Field, M., & Jones, A. (2018). Reprint of" Methodological and reporting quality in laboratory studies of human eating behavior". *Appetite*, 130, 321-326.
- Rogers, P. J. & Hill, A. J. (1989) Breakdown of dietary restraint following mere exposure to food stimuli: Interrelationships between restraint, hunger, salivation, and food intake. *Addictive behaviors*, 14(4), 387-197.
- Schifferstein, H. N., Fenko, A., Desmet, P. M., Labbe, D., & Martin, N. (2013). Influence of package design on the dynamics of multisensory and emotional food experience. *Food Quality and Preference*, 27(1), 18-25.
- Schubert, T., Friedmann, F., & Regenbrecht, H. (2001). The experience of presence: Factor analytic insights. *Presence: Teleoperators & Virtual Environments*, *10*(3), 266-281.
- Sharma, V., & Estes, Z. (2024). Seeing is smelling: Pictures improve product evaluations by evoking olfactory imagery. *International Journal of Research in Marketing*, 41(2), 282-307.
- Ship, J. A., Fox, P. C., & Baum, B. J. (1991). How much saliva is enough?. *The Journal of the American Dental Association*, *122*(3), 63-69.
- Slater, M. (2018) Immersion and the illusion of presence in virtual reality. *British Journal of Psychology*, 109, 431-433.

Slater, M., & Wilbur, S. (1997). A Framework for Immersive Virtual Environments (FIVE):

Speculations on the Role of Presence in Virtual Environments. *Presence: Teleoperators and Virtual Environments*, 6(6), 603–616. https://doi.org/10.1162/pres.1997.6.6.603

- Smeets, M. A., Schifferstein, H. N., Boelema, S. R., & Lensvelt-Mulders, G. (2008). The Odor Awareness Scale: A new scale for measuring positive and negative odor awareness. *Chemical senses*, 33(8), 725-734.
- Spence, C. (2011). Mouth-watering: the influence of environmental and cognitive factors on salivation and gustatory/flavor perception. *Journal of Texture Studies*, *42*(2), 157-171.
- Steel, D., Kemps, E., & Tiggemann, M. (2006). Effects of hunger and visuo-spatial interference on imagery-induced food cravings. *Appetite*, *46*(1), 36-40.
- Tiggemann, M., & Kemps, E. (2005). The phenomenology of food cravings: The role of mental imagery. *Appetite*, *45*(3), 305-313.
- Tuorila, H., Lähteenmäki, L., Pohjalainen, L., & Lotti, L. (2001). Food neophobia among the Finns and related responses to familiar and unfamiliar foods. *Food Quality and Preference*, 12(1), 29-37.
- de Vries, R., Boesveldt, S., & de Vet, E. (2021). Locating calories: Does the high-calorie bias in human spatial memory influence how we navigate the modern food environment?.
 Food Quality and Preference, 94, 104338.
- van der Waal, N. E., Janssen, L., Antheunis, M., Culleton, E., & van der Laan, L. N. (2021).
 The appeal of virtual chocolate: A systematic comparison of psychological and physiological food cue responses to virtual and real food. *Food Quality and Preference*, *90*, 104167.
- Waterlander, W.E., Jiang, Y., Steenhuis, I. H. M., Mhurchu, C. N. (2015) Using a 3D virtual supermarket to measure food purchase behavior: a validation study. *Journal of Medical Internet Research*, 17, e107.

- White, K. D. (1978). Salivation: The significance of imagery in its voluntary control. *Psychophysiology*, *15*(3), 196-203.
- Xu, C., Siegrist, M., Hartmann, C. (2021) The application of virtual reality in food consumer behavior research: a systematic review. *Trends in Food Science & Technology*, 116, 533-544.
- Yang, X., Zandstra, E. H., & Boesveldt, S. (2023). How sweet odors affect healthy food choice: An eye-tracking study. *Food Quality and Preference*, 109, 104922. https://doi.org/10.1016/j.foodqual.2023.104922
- Zeman, A., Milton, F., Della Sala, S., Dewar, M., Frayling, T., Gaddum, J., ... & Winlove, C. (2020). Phantasia–the psychological significance of lifelong visual imagery vividness extremes. *Cortex*, 130, 426-440.